

Search  for

☒ Limits
 [Preview/Index](#)
[History](#)
[Clipboard](#)
[Details](#)

[About Entrez](#)

[Text Version](#)

Items 1-20 of 164

of 9
 [Next](#)

#### Entrez PubMed

[Overview](#)  
[Help | FAQ](#)  
[Tutorial](#)  
[New/Noteworthy](#)  
[E-Utilities](#)

#### PubMed Services


[Journals Database](#)  
[MeSH Database](#)  
[Single Citation Matcher](#)  
[Batch Citation Matcher](#)  
[Clinical Queries](#)  
[LinkOut](#)  
[Cubby](#)

#### Related Resources


[Order Documents](#)  
[NLM Gateway](#)  
[TOXNET](#)  
[Consumer Health](#)  
[Clinical Alerts](#)  
[ClinicalTrials.gov](#)  
[PubMed Central](#)

[Privacy Policy](#)


☐ **1:** [Levitan D, Greenwald I.](#) [Related Articles, Links](#)


**Facilitation of lin-12-mediated signalling by sel-12, a *Caenorhabditis elegans* S182 Alzheimer's disease gene.**  
 Nature. 1995 Sep 28;377(6547):351-4.  
 PMID: 7566091 [PubMed - indexed for MEDLINE]


☐ **2:** [Grant B, Greenwald I.](#) [Related Articles, Links](#)


**The *Caenorhabditis elegans* sel-1 gene, a negative regulator of lin-12 and glp-1, encodes a predicted extracellular protein.**  
 Genetics. 1996 May;143(1):237-47.  
 PMID: 8722778 [PubMed - indexed for MEDLINE]


☐ **3:** [Hubbard EJ, Wu G, Kitajewski J, Greenwald I.](#) [Related Articles, Links](#)


**sel-10, a negative regulator of lin-12 activity in *Caenorhabditis elegans*, encodes a member of the CDC4 family of proteins.**  
 Genes Dev. 1997 Dec 1;11(23):3182-93.  
 PMID: 9389650 [PubMed - indexed for MEDLINE]


☐ **4:** [Doyle TG, Wen C, Greenwald I.](#) [Related Articles, Links](#)


**SEL-8, a nuclear protein required for LIN-12 and GLP-1 signaling in *Caenorhabditis elegans*.**  
 Proc Natl Acad Sci U S A. 2000 Jul 5;97(14):7877-81.  
 PMID: 10884418 [PubMed - indexed for MEDLINE]


☐ **5:** [Wen C, Greenwald I.](#) [Related Articles, Links](#)


**p24 proteins and quality control of LIN-12 and GLP-1 trafficking in *Caenorhabditis elegans*.**  
 J Cell Biol. 1999 Jun 14;145(6):1165-75.  
 PMID: 10366590 [PubMed - indexed for MEDLINE]


☐ **6:** [Hubbard EJ, Dong Q, Greenwald I.](#) [Related Articles, Links](#)












**Evidence for physical and functional association between EMB-5 and LIN-12 in *Caenorhabditis elegans*.**  
 Science. 1996 Jul 5;273(5271):112-5.  
 PMID: 8658178 [PubMed - indexed for MEDLINE]

☐ **7:** [Wen C, Metzstein MM, Greenwald I.](#) [Related Articles, Links](#)


**SUP-17, a *Caenorhabditis elegans* ADAM protein related to *Drosophila* KUZBANIAN, and its role in LIN-12/NOTCH signalling.**  
 Development. 1997 Dec;124(23):4759-67.  
 PMID: 9428412 [PubMed - indexed for MEDLINE]

☐ **8:** [Li X, Greenwald I.](#) [Related Articles, Links](#)


**HOP-1, a *Caenorhabditis elegans* presenilin, appears to be functionally redundant with SEL-12 presenilin and to facilitate LIN-12 and GLP-1 signaling.**  
 Proc Natl Acad Sci U S A. 1997 Oct 28;94(22):12204-9.  
 PMID: 9342387 [PubMed - indexed for MEDLINE]

- ☐ **9:** [Tax FE, Thomas JH, Ferguson EL, Horvitz HR.](#) [Related Articles, Links](#)  
 **Identification and characterization of genes that interact with lin-12 in *Caenorhabditis elegans*.**  
Genetics. 1997 Dec;147(4):1675-95.  
PMID: 9409830 [PubMed - indexed for MEDLINE]
- ☐ **10:** [Westlund B, Parry D, Clover R, Basson M, Johnson CD.](#) [Related Articles, Links](#)  
 **Reverse genetic analysis of *Caenorhabditis elegans* presenilins reveals redundant but unequal roles for sel-12 and hop-1 in Notch-pathway signaling.**  
Proc Natl Acad Sci U S A. 1999 Mar 2;96(5):2497-502.  
PMID: 10051671 [PubMed - indexed for MEDLINE]
- ☐ **11:** [Goutte C, Hepler W, Mickey KM, Priess JR.](#) [Related Articles, Links](#)  
 **aph-2 encodes a novel extracellular protein required for GLP-1-mediated signaling.**  
Development. 2000 Jun;127(11):2481-92.  
PMID: 10804188 [PubMed - indexed for MEDLINE]
- ☐ **12:** [Sundaram M, Greenwald I.](#) [Related Articles, Links](#)  
 **Suppressors of a lin-12 hypomorph define genes that interact with both lin-12 and glp-1 in *Caenorhabditis elegans*.**  
Genetics. 1993 Nov;135(3):765-83.  
PMID: 8293978 [PubMed - indexed for MEDLINE]
- ☐ **13:** [Levitan D, Greenwald I.](#) [Related Articles, Links](#)  
 **Effects of SEL-12 presenilin on LIN-12 localization and function in *Caenorhabditis elegans*.**  
Development. 1998 Sep;125(18):3599-606.  
PMID: 9716525 [PubMed - indexed for MEDLINE]
- ☐ **14:** [Levitan D, Yu G, St George Hyslop P, Goutte C.](#) [Related Articles, Links](#)  
 **APH-2/nicastrin functions in LIN-12/Notch signaling in the *Caenorhabditis elegans* somatic gonad.**  
Dev Biol. 2001 Dec 15;240(2):654-61.  
PMID: 11784090 [PubMed - indexed for MEDLINE]
- ☐ **15:** [Christensen S, Kodoyianni V, Bosenberg M, Friedman L, Kimble J.](#) [Related Articles, Links](#)  
 **lag-1, a gene required for lin-12 and glp-1 signaling in *Caenorhabditis elegans*, is homologous to human CBF1 and *Drosophila* Su(H).**  
Development. 1996 May;122(5):1373-83.  
PMID: 8625826 [PubMed - indexed for MEDLINE]
- ☐ **16:** [Baumeister R, Leimer U, Zweckbronner I, Jakubek C, Grunberg J, Haass C.](#) [Related Articles, Links](#)  
 **Human presenilin-1, but not familial Alzheimer's disease (FAD) mutants, facilitate *Caenorhabditis elegans* Notch signalling independently of proteolytic processing.**  
Genes Funct. 1997 Apr;1(2):149-59.  
PMID: 9680315 [PubMed - indexed for MEDLINE]
- ☐ **17:** [Fares H, Greenwald I.](#) [Related Articles, Links](#)  
 **SEL-5, a serine/threonine kinase that facilitates lin-12 activity in *Caenorhabditis elegans*.**  
Genetics. 1999 Dec;153(4):1641-54.  
PMID: 10581273 [PubMed - indexed for MEDLINE]
- ☐ **18:** [Cinar HN, Sweet KL, Hosemann KE, Earley K, Newman AP.](#) [Related Articles, Links](#)  
 **The SEL-12 presenilin mediates induction of the *Caenorhabditis elegans* uterine pi cell fate.**  
Dev Biol. 2001 Sep 1;237(1):173-82.

☐ **19:** [Eimer S, Lakowski B, Donhauser R, Baumeister R.](#)

[Related Articles, Links](#)



Loss of spr-5 bypasses the requirement for the C.elegans presenilin sel-12 by derepressing hop-1.

EMBO J. 2002 Nov 1;21(21):5787-96.

PMID: 12411496 [PubMed - indexed for MEDLINE]

☐ **20:** [Hoskins R, Hajnal AF, Harp SA, Kim SK.](#)

[Related Articles, Links](#)



The C. elegans vulval induction gene lin-2 encodes a member of the MAGUK family of cell junction proteins.

Development. 1996 Jan;122(1):97-111.

PMID: 8565857 [PubMed - indexed for MEDLINE]

Display	Summary	Show: 20	Sort	Send to	Text
Items 1-20 of 164				Page 1	of 9 Next

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)